



King's Research Portal

DOI:

[10.1136/gutjnl-2018-316225](https://doi.org/10.1136/gutjnl-2018-316225)

Document Version

Version created as part of publication process; publisher's layout; not normally made publicly available

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Gudra, D., Shoaie, S., Fridmanis, D., Klovins, J., Wefer, H., Silamikelis, I., Peculis, R., Kalnina, I., Elbere, I., Radovica-Spalvina, I., Hultcrantz, R., Senders, ., Leja, M., & Engstrand, L. (2018). A widely used sampling device in colorectal cancer screening programmes allows for large-scale microbiome studies. *Gut*. <https://doi.org/10.1136/gutjnl-2018-316225>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

LETTER

A widely used sampling device in colorectal cancer screening programmes allows for large-scale microbiome studies

We read with interest the article by Passamonti *et al.*,¹ reporting the performance of two different faecal immunochemical tests (FITs) highlighting the importance of standardisation and validation of screening methodologies. Conventionally, laboratory-based FIT is the preferred approach in testing for occult blood in faeces, which includes colorectal cancer screening programmes.²⁻⁴ The potential of preserving stable faecal samples in a widely used FIT buffer for microbiome research would enable prospective microbiome studies in generally healthy subjects undergoing colorectal cancer screening.

For this purpose, we evaluated faecal sample stability in the commonly used OC-Sensor (Eiken Chemical, Tokyo, Japan) under various storage conditions. Faecal samples from five healthy adult individuals were used for the analysis and exposed to 16 different conditions: immediately frozen at -86°C (with FIT (wFIT)/without FIT (woFIT)); immediately frozen at -20°C (wFIT/woFIT); wFIT stored at

4°C for 1, 2, 7, 14 days; stored at 20°C for 1, 2, 7, 14 days; stored at 30°C for 1, 2, 7 days and lastly stored at 4°C for 2 days and at 20°C for additional 2 days. Shotgun metagenomic analysis was performed by Illumina 2500, while taxonomic compositions were determined by 16S rRNA analysis employing Illumina MiSeq and Ion Torrent PGM.

Our first attempt to extract DNA from wFIT samples failed due to low amount of DNA obtained. Thus, lyophilisation for all of the wFIT samples, except for woFIT samples, was applied, increasing the yield of DNA up to 30 times.

As shown by the figure 1, the Shannon index in both platforms displayed a clear pattern of decreased diversity during prolonged storage. While testing for differences between samples we discovered that woFIT samples, wFIT samples stored at 4°C for 1 and 2 days, wFIT samples stored at 20°C for 1 day from the Illumina MiSeq data significantly differ from wFIT samples stored at 30°C for 7 days (one-way analysis of variance; $p < 0.05$).

The similarity matrix using unweighted principal coordinate analysis (PCoA, figure 2A,C) demonstrated that samples collected from each individual clustered together and was consistent in data gained from both sequencing platforms, suggesting that the biological effect outweighed the effect of the sample handling. Further, the weighted PCoA

(figure 2B,D) displays that native samples tend to cluster closer while samples stored for prolonged periods tend to be located further away from the native ones.

Comparing observed operational taxonomic units (OTUs) between samples that were stored within various storage conditions, we were unable to identify any specific OTU cluster across all taxonomical levels that would significantly differ (Kruskal-Wallis test) between any storage condition. Nevertheless, we found that the relative abundances of Gram-negative bacteria tended to decrease while the abundances for most of the Gram-positive bacteria tended to increase over time. However, it is possible that even within native samples certain bacterial OTUs could fluctuate regardless of the sequencing platform employed.

Shotgun sequencing was performed on 12 samples (figure 2E) from four individuals: six woFIT samples (WO1-4) that were immediately stored at -86°C and six wFIT samples from three individuals (FIT1-4), stored for 2 days at 4°C and for 2 days at 20°C to imitate the typical sample treatment procedure when the material is being transported to a central laboratory by standard mail delivery. In addition to access the reproducibility of acquired data, technical replicates in the form of DNA that was extracted from independently collected samples from individuals 2 and 3 were included within




Figure 1 Mean Shannon diversity index values by storage conditions of faecal samples sequenced on Ion Torrent Personal Genome Machine and Illumina MiSeq. Error bars represent SEs. FIT, faecal immunochemical test.




Figure 2 Beta diversity (panels A–D) analysis of GI tract bacterial communities between the individuals and storage conditions presented in a form of principal coordinate analysis (PCoA) plot of weighted and unweighted UniFrac distances. Panels A and C are coloured according to the individual, panels B and D are coloured according to the conditions that samples were exposed to. Panels A and B represent the UniFrac metrics obtained from sequencing on the Illumina MiSeq platform, whereas panels C and D represent UniFrac metrics obtained from sequencing on the Ion Torrent PGM platform. The results of the metagenomics analysis at the species level are presented within panel E in a form of the heat-map profile for the most abundant entities in the metagenome samples. According to upper dendrogram, there is a consistency within the samples composition as the samples from the same individual are clustering together. Samples WO1, WO2.1, WO2.2, WO3.1, WO3.2, WO4 were without faecal immunochemical test (woFIT) and immediately frozen at -86°C ; while samples FIT1, FIT2.1, FIT2.2, FIT3.1, FIT3.2 and FIT4 were with FIT (wFIT) and stored at 4°C for 2 days and then at 20°C for additional 2 days. Samples FIT2.2 and FIT4 failed at the sequencing stage and therefore are not presented within the figure. Both sample groups include technical replicates for individuals 2 and 3 (eg, WO2.1. and WO2.2.; WO3.1 and WO3.2.; FIT2.1 and FIT2.2.; FIT3.1. and FIT3.2.).

this analysis. However, two wFIT samples (eg, FIT2.2 and FIT4) failed during the sequencing stage and therefore were excluded from further analysis. A similarity matrix was built with relative abundances to calculate the PCoA. Figure 2E illustrates the similarity between species for two types of sampling showing that results are not biased to the sampling procedures. Results indicate that sample similarities tended towards samples from the same subject rather than the storage method.

In this study, we sequenced microbiome samples employing two different sequencing platforms—Ion Torrent PGM that allows to analyse one variable region (V3) and to verify acquired results and also Illumina MiSeq platform that enables the analysis of two variable regions (V3–V4) at a time. Analysing absolute OTU frequencies within the samples, we did not find a significant difference between immediately frozen samples, which coincide with previous studies.^{5–8} Although previous study⁹ has found lower FIT stability measures, we observed some limitations as they extracted DNA from a

small proportion of sample, while in our study design lyophilisation procedure was included allowing to extract DNA from whole volume. However, we did not include technical replicates for 16S rRNA analysis. Another critical aspect in FIT-oriented studies is the variety of available FIT test tubes in the market. Each FIT buffer from various manufacturers might contain slightly different ingredients and concentrations,¹⁰ which seems to be a trade secret and might alter the microbial composition. In our study, employed OC-Sensor is a reliable and convenient sampling device that can be used in large-scale microbiome studies on a population level even outside colorectal cancer screening programmes.

Dita Gudra,¹ Saeed Shoaie,^{2,3} Davids Fridmanis,¹ Janis Klovins,¹ Hugo Wefer,^{2,4} Ivars Silamikelis,¹ Raitis Peculis,¹ Ineta Kalnina,¹ Ilze Elbere,¹ Ilze Radovica-Spalvina,¹ Rolf Hultcrantz,² Ģirts Šķenders,^{5,6} Marcis Leja,^{5,6} Lars Engstrand^{2,4}

¹Latvian Biomedical Research and Study Centre, Riga, Latvia

²Centre for Translational Microbiome Research, Department of Microbiology Tumor and Cell Biology, Karolinska Institutet, Stockholm, Sweden

³Centre for Host–Microbiome Interactions, Dental Institute, King's College London, London, UK

⁴Science for Life Laboratory, Solna, Sweden

⁵Faculty of Medicine, University of Latvia, Riga, Latvia

⁶Institute of Clinical and Preventive Medicine, University of Latvia, Riga, Latvia

Correspondence to Dr Davids Fridmanis, Latvian Biomedical Research and Study Centre, Ratsupites 1K-1, LV-1067, Riga, Latvia; davids@biomed.lu.lv.
Iv and Professor Lars Engstrand, Department of Microbiology, Tumor and Cell Biology, Karolinska Institutet, Stockholm 171 77, Sweden; lars.engstrand@ki.se

Contributors Study concept and design was created by ML and LE. Data collection was performed by DG, HW, IR-S, RH, IK, IE, GS. Bioinformatics analysis performed by DG, SS, IS, RP. DG and SS wrote the manuscript. Funding was obtained by LE, DF, JK and ML. All authors approved the final version for publication.

Funding The work was supported by the ERDF project 'Long-term effects of H. pylori eradication on gastrointestinal tract microbiome and development of screening system for detection of extended-spectrum beta-lactamase coding genes within faeces samples' (project no: 1.1.1.1/16/A/272) and in part by the Swedish Research Council (project no: K2013-56X-10848) and the Söderbergs Foundation.

Competing interests None declared.

Patient consent Obtained.

Ethics approval Biomedical Ethics Committee of the Riga East University Hospital Support Foundation, approval no: 13-A/13.

Provenance and peer review Not commissioned; externally peer reviewed.



OPEN ACCESS

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

© Author(s) (or their employer(s)) 2018. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

ML and LE contributed equally.
DG and SS contributed equally.



To cite Gudra D, Shoaie S, Fridmanis D, *et al.* *Gut* Epub ahead of print: [please include Day Month Year]. doi:10.1136/gutjnl-2018-316225

Received 9 February 2018

Revised 8 August 2018

Accepted 26 August 2018

Gut 2018;0:1–3. doi:10.1136/gutjnl-2018-316225

REFERENCES

- 1 Passamonti B, Malaspina M, Fraser CG, *et al.* A comparative effectiveness trial of two faecal immunochemical tests for haemoglobin (FIT). Assessment of test performance and adherence in a single round of a population-based screening programme for colorectal cancer. *Gut* 2018;67:485–96.
- 2 Directorate-General for Health and Consumers (European Commission) EAfHaC, World Health Organization. *European guidelines for quality assurance in colorectal cancer screening and diagnosis*. Luxembourg, 2010.
- 3 Young GP, Symonds EL, Allison JE, *et al.* Advances in fecal occult blood tests: the FIT revolution. *Dig Dis Sci* 2015;60:609–22.
- 4 Robertson DJ, Lee JK, Boland CR, *et al.* Recommendations on fecal immunochemical testing to screen for colorectal neoplasia: a consensus statement by the us multi-society task force on colorectal cancer. *Am J Gastroenterol* 2017;112:37–53.
- 5 Hale VL, Tan CL, Knight R, *et al.* Effect of preservation method on spider monkey (*Ateles geoffroyi*) fecal microbiota over 8 weeks. *J Microbiol Methods* 2015;113:16–26.
- 6 Carroll IM, Ringel-Kulka T, Siddle JP, *et al.* Characterization of the fecal microbiota using high-throughput sequencing reveals a stable microbial community during storage. *PLoS One* 2012;7:e46953.
- 7 Vlčková K, Mrázek J, Kopečný J, *et al.* Evaluation of different storage methods to characterize the fecal bacterial communities of captive western lowland gorillas (*Gorilla gorilla gorilla*). *J Microbiol Methods* 2012;91:45–51.
- 8 Cardona S, Eck A, Cassellas M, *et al.* Storage conditions of intestinal microbiota matter in metagenomic analysis. *BMC Microbiol* 2012;12:158.
- 9 Vogtmann E, Chen J, Kibriya MG, *et al.* Comparison of fecal collection methods for microbiota studies in Bangladesh. *Appl Environ Microbiol* 2017;83:e00361–17.
- 10 Grow MA, Shah VD. *Fecal sample immunoassay method testing for hemoglobin*. European Patent Office., 1993.